

# Dislocation structure and kinetics in slip-twin model



Max-Planck-Institut  
für Eisenforschung GmbH

MSU Twin Meeting, Duesseldorf – October 6<sup>th</sup>, 2009



## Internal variables:

- $N^\alpha$  edge dislocation densities  $\varrho_{\text{edge}}^\alpha$
- $N^\alpha$  dipole densities  $\varrho_{\text{dipole}}^\alpha$

## Derived measures:

- $\tau_c^\alpha$  threshold shear stress
- $\lambda^\alpha$  mean distance between 2 obstacles seen by a dislocation

Threshold stress  $\tau_c^\alpha$ :

$$\tau_c^\alpha = G_{\text{iso}} b^\alpha \sqrt{\sum_{\tilde{\alpha}=1}^{N^\alpha} \xi^{\alpha\tilde{\alpha}} (\varrho_{\text{edge}}^{\tilde{\alpha}} + \varrho_{\text{dipole}}^{\tilde{\alpha}})}$$

with:

- $G_{\text{iso}}$  Isotropic shear modulus
- $b^\alpha$  Burgers vector of slip system  $\alpha$
- $\xi^{\alpha\tilde{\alpha}}$  interaction strength (Kubin et al. 2008)



Shear rate  $\dot{\gamma}^\alpha$ :

$$\dot{\gamma}^\alpha = \rho_{\text{edge}}^\alpha b^\alpha v_{\text{glide}}^\alpha$$

Velocity  $v_{\text{glide}}^\alpha$ :

$$v_{\text{glide}}^\alpha = v_0 \exp \left[ -\frac{Q}{k_B T} \left( 1 - \left( \frac{|\tau^\alpha|}{\tau_c^\alpha} \right)^p \right)^q \right] \text{sign}(\tau^\alpha)$$

with:

- $v_0$  Velocity pre-factor
- $Q$  Activation energy for dislocation glide
- $k_B T$  Boltzmann energy



Multiplication:

$$\dot{\rho}_{\text{multiplication}}^{\alpha} = \frac{|\dot{\gamma}^{\alpha}|}{b^{\alpha} \lambda^{\alpha}}$$



Dipole formation:

$$\dot{\varrho}_{\text{formation}}^{\alpha} = 2 \frac{\hat{d}^{\alpha}}{b^{\alpha}} \frac{\varrho_{\text{edge}}^{\alpha}}{2} |\dot{\gamma}^{\alpha}|$$

Length  $\hat{d}^{\alpha}$ :

$$\hat{d}^{\alpha} = \frac{1}{8\pi} \frac{G_{\text{iso}} b^{\alpha}}{1 - \nu} \frac{1}{|\tau^{\alpha}|}$$



Single-single annihilation:

$$\dot{\rho}_{\text{single-single}}^{\alpha} = 2 \frac{2 \check{d}^{\alpha}}{b^{\alpha}} \frac{\rho_{\text{edge}}^{\alpha}}{2} |\dot{\gamma}^{\alpha}|$$

Length  $\check{d}^{\alpha}$ :

$$\check{d}^{\alpha} \propto b^{\alpha}$$

# Spontaneous annihilation of one single dislocation and one dipole constituent



Single-dipole constituent annihilation:

$$\dot{\varrho}_{\text{single-dipole}}^{\alpha} = 2 \frac{2 \check{d}^{\alpha}}{b^{\alpha}} \frac{\varrho_{\text{dipole}}^{\alpha}}{2} |\dot{\gamma}^{\alpha}|$$



Dipole climb:

$$\dot{\varrho}_{\text{climb}}^{\alpha} = \varrho_{\text{dipole}}^{\alpha} \frac{4 v_{\text{climb}}}{\hat{d}^{\alpha} + \check{d}^{\alpha}}$$

Climb velocity  $v_{\text{climb}}^{\alpha}$ :

$$v_{\text{climb}}^{\alpha} = \frac{D \Omega^{\alpha}}{b^{\alpha} k_B T} \frac{G_{\text{iso}} b^{\alpha}}{2 \pi (1 - \nu)} \frac{2}{\hat{d}^{\alpha} + \check{d}^{\alpha}}$$



Edge dislocation density rate:

$$\dot{\rho}_{\text{edge}}^{\alpha} = \dot{\rho}_{\text{multiplication}}^{\alpha} - \dot{\rho}_{\text{formation}}^{\alpha} - \dot{\rho}_{\text{single-single}}^{\alpha}$$

Dislocation dipole density rate:

$$\dot{\rho}_{\text{dipole}}^{\alpha} = \dot{\rho}_{\text{formation}}^{\alpha} - \dot{\rho}_{\text{single-dipole}}^{\alpha} - \dot{\rho}_{\text{climb}}^{\alpha}$$